















R4D Community Types or Subtypes	Yukon Delta	Chukchi-Imuruk	Cape Thompson	NW Alaska	Atkasook	Umiat	Arrigetch Peaks	Yukon	Yukon, N.W.T.	Alaska Statewide Classification	Alaska Statewide Classification
Walker (1987, Level C)	Tande and Jennings (1986)	Racine and Anderson (1979)	Johnson et al. (1966)	Hanson (1953)	Komarkova and Webber (1980)	Churchill (1955)	Cooper (1986)	Lambert (1968)	Ritchie (1984)	Viereck et al. Level IV (1986)	Viereck et al. Level V (1986)
Aquatic <i>Arctophila fulva</i> grass marsh	Wet <i>Arctophila fulva</i> -( <i>Hippuris tetraphylla</i> ) grass forb type (W5a)	Aquatic meadow	Aquatic community		<i>Arctophila fulva</i> seasonal short grass			<i>Arctophiletum fulvae</i>		Fresh grass marsh [3.A.(3) e.]	<i>Arctophila fulva</i> (Batten 1977; Bergman et al. 1977; Britton 1967; Childs 1969; Clebsch 1957; Hulten 1966; Komarkova and Webber 1978; Murray 1974; Potter 1972; Racine and Anderson 1979; Rausch and Rausch 1968; Strevler et al. 1973; Webber et al. 1978; Wiggins and Thomas 1962)
Dry <i>Rhizocarpon geographicum</i> , <i>Cetraria nigricans</i> lichen barren		Rock desert		Lichen-moss barrens			<i>Umbilicarietum-pensylvanico-caroliniana</i>			Crustose lichen [3.C.(2)a.]	<i>Umbilicaria</i> spp.- <i>Rhizocarpon</i> spp. (Anderson 1974; Hanson 1953; Kessel and Schaller 1960; Klein 1958; Pegau 1968; Rausch and Rausch 1968; Webber et al. 1978)
Dry <i>Festuca rubra</i> , <i>Poa glauca</i> grass barren				Grassland type						Midgrass herb [3.A.(1)d.]	